

Magnetic field control of elastic scattering in a cold gas of fermionic lithium atoms

S. Jochim, M. Bartenstein, G. Hendl, J. Hecker Denschlag, and R. Grimm

Institut für Experimentalphysik, Universität Innsbruck, Technikerstr. 25, 6020 Innsbruck, Austria

A. Mosk

FOM instituut voor plasmafysica Rijnhuizen, P.O. Box 1207, 3430 BE Nieuwegein, The Netherlands

M. Weidemüller

Max-Planck-Institut für Kernphysik, Postfach 103980, 69029 Heidelberg, Germany

(Dated: July 24, 2002)

We study elastic collisions in an optically trapped spin mixture of fermionic lithium atoms in the presence of magnetic fields up to 1.5 kG by measuring evaporative loss. Our experiments confirm the expected magnetic tunability of the scattering length by showing the main features of elastic scattering according to recent calculations. We measure the zero crossing of the scattering length that is associated with a predicted Feshbach resonance at 530(3) G. Beyond the resonance we observe the expected large cross section in the triplet scattering regime.

PACS numbers: 34.50.-s, 05.30.Fk, 39.25.+k, 32.80.Pj

In an ultracold atomic gas, the s -wave scattering length a characterizes the elastic interactions and is therefore an especially interesting parameter to manipulate. Changes of a have a profound effect on the dynamics of the gas and on its thermodynamic stability. For spin mixtures of fermionic atoms, the scattering length governs the stability and critical temperature of a Cooper-paired superfluid phase. The scattering length can be conveniently tuned if it depends on the magnetic field, as is the case near a Feshbach resonance. For bosonic atoms, such resonances have been observed [1, 2, 3], and used for attainment and manipulation of a BEC in ^{85}Rb [4] and for the production of bright solitons in bosonic ^7Li [5, 6].

For fermionic gases, Loftus *et al.* have recently observed a Feshbach resonance between two different spin states of ^{40}K , which changes scattering properties in a narrow magnetic field range [7]. The other fermionic species currently used in several experiments, ^6Li , is predicted to have a very broad Feshbach resonance which completely changes the character of interactions in a wide experimentally accessible field range. Magnetic-field tuning of elastic scattering at small fields far below the resonance was recently used by Granade *et al.* to obtain a sufficient scattering cross section for the all-optical production of a degenerate Fermi gas of lithium [8]. At higher magnetic fields, an expected extreme tunability promises great prospects to observe new phenomena, e.g. in Cooper pairing [9, 10], cold molecule formation [11] and studies of Fermi gases in reduced dimensionality [12].

In this Letter, we experimentally explore the magnetic tunability of elastic scattering in a spin-mixture of fermionic lithium atoms in high magnetic fields up to 1.5 kG. Our results verify the expected dependence of the cross section that follows from theoretical calculations of the scattering length [13, 14, 15]. As a particular feature associated with a predicted Feshbach resonance [13], we




FIG. 1: (a) Model curve approximating the results of [13, 14, 15] for the s -wave scattering length of ^6Li atoms in the two lowest spin states versus magnetic field. (b) Corresponding behavior of the scattering cross section at a finite collision energy with a relative wave number of $k = (300a_0)^{-1}$.

observe the zero crossing of the scattering length that occurs at a field of 530 G. The exact location of this feature is of great interest as a sensitive input parameter to better constrain the uncertainty in the molecular potentials for more accurate theoretical calculations of the scattering properties of ^6Li . Our measurements of elastic collisions are based on evaporation out of an optical dipole trap. We use an optical standing-wave trap developed earlier [16], which employs the power enhancement inside an optical resonator to create a deep trap with a sufficiently large volume to capture a large number of lithium atoms.

The scattering properties in different spin-mixtures of fermionic lithium atoms were theoretically investigated by Houbiers *et al.* [13], Kokkelmans *et al.* [14], and Ven-

turi and Williams [15]. Magnetic tunability, of particular interest for Cooper pairing in a Fermi gas [9, 10], was predicted for the stable combination of the two lowest states $|1\rangle$ and $|2\rangle$; at low magnetic field these states correspond to $F = 1/2$, $m_F = +1/2$ and $m_F = -1/2$, respectively. Most prominently, a broad Feshbach resonance is expected to mark the transition from the low-field scattering regime to the high-field region. In more detail, the s -wave scattering length $a(B)$ is expected to show the following behavior: After being zero in the absence of a magnetic field, a local maximum of $|a(B)|$ occurs with rising field at ~ 300 G with $a \approx -300 a_0$, where a_0 is the Bohr radius. Then, as a precursor of the Feshbach resonance, the scattering length crosses zero in the range between 500 G and 550 G. After passing through the resonance somewhere between 800 G and 850 G scattering in higher fields is dominated by the triplet potential with a very large and negative scattering length. The available theoretical data [13, 14, 15] show the same behavior for $a(B)$ within some variations due to the limited knowledge of the molecular interaction parameters. Fig. 1(a) illustrates these predictions for the scattering length $a(B)$ by a corresponding model curve [21] that approximates the results of Refs. [13, 14, 15].

In a cold gas at finite temperature the cross section for elastic scattering of non-identical particles is unitarity-limited to a maximum value of $\sigma_{\max} = 4\pi/k^2$, where $k = mv/(2\hbar)$ is the wavenumber corresponding to a relative velocity v and a reduced mass $m/2$. Taking into account the B-field dependent scattering length $a(B)$ and the unitarity limit, the resulting B-field dependent cross section can be written as $\sigma = 4\pi a^2/(1 + k^2 a^2)$. For the considered $|1\rangle - |2\rangle$ spin mixture of ^6Li the expected behavior of the cross section is shown in Fig. 1(b) for the example of a wave number $k = (300 a_0)^{-1}$ close to our experimental conditions. Most notably, as a consequence of the unitarity limit in combination with the very large scattering length for high magnetic fields, the Feshbach resonance does not appear as a pronounced feature in the cross section. The zero crossing of the scattering length, however, leads to a vanishing scattering cross section and thus shows up as a manifestation of the resonance.

The resonator-enhanced dipole trap (REDT) [16] makes use of the enhancement of the laser intensity inside a linear optical resonator to create a large and deep trapping volume for lithium atoms. The power provided by a 2-W Nd:YAG laser (Innolight Mephisto-2000) at a wavelength of 1064 nm is enhanced by a factor of 120 to create a far red-detuned 1D optical lattice trap with an axial period of 532 nm and a transverse $1/e$ -radius of $115 \mu\text{m}$. The maximum trap depth is of the order of 1 mK. To vary the trap depth the resonator-internal power is servo-controlled by an acousto-optical modulator in the incident laser beam. From a standard magneto-optical trap (MOT) operated with diode lasers we typically transfer 5×10^5 ^6Li atoms into roughly 1000 individual wells at




FIG. 2: Evaporative loss measurements over the full magnetic field range. The data points show the measured number of atoms remaining in the trap after 1 s (○) and 3 s (●) of plain evaporation at a constant trap depth of $750 \mu\text{K}$.

a temperature of $\sim 400 \mu\text{K}$. The resulting peak density is $\sim 1.5 \times 10^{11} \text{ cm}^{-3}$. By extinguishing the repumping light of the MOT 1 ms before the main trapping light is turned off, all atoms are pumped into the two states $|1\rangle$ and $|2\rangle$ to create the desired spin mixture.

The magnetic field is produced by a pair of water-cooled coils outside of the glass vacuum cell of the trap. At a maximum continuous operation current of 200 A the coils produce a magnetic field of 1.5 kG with a curvature of only 75 G/cm^2 along the symmetry axis; the corresponding power dissipation is 6 kW. The set-up allows for a maximum ramp speed of 5 G/ms within the full range. The magnetic field is calibrated by radio-frequency induced transitions from $|2\rangle$ to the state that at $B = 0$ corresponds to $F = 3/2$, $m_F = +1/2$. The latter is unstable against inelastic collisions with $|2\rangle$ which leads to easily detectable loss. With a fit to the Breit-Rabi formula we obtain a calibration of the magnetic field to better than 1 G over the full range.

The basic idea of our measurements is to observe elastic collisions through evaporative loss at a variable magnetic field [17]. Our trap is initially loaded under conditions where the effective temperature T of a truncated Boltzmann distribution [18] is only slightly below the trap depth U . A strongly non-thermal distribution is thus created with a small truncation parameter $\eta = U/k_B T \approx 2$. The thermal relaxation resulting from elastic collisions then leads to rapid evaporative loss and cooling of the sample, i.e. an increase of η . The trap depth can be kept constant to study plain evaporation or, alternatively, ramped down to force the evaporation process.

In a series of plain evaporation experiments performed at a constant trap depth of $750 \mu\text{K}$ we measure evaporative loss over the maximum accessible range of magnetic fields up to 1.5 kG. After a fixed holding time the remain-

ing atoms are retrapped into the MOT and their number is measured via the fluorescence signal by a calibrated photodiode. The loss signal is recorded for holding times of 1 s and 3 s corresponding to the time scale of evaporation. These holding times are short compared with the rest-gas limited lifetime of 30 s. Fig. 2 shows the result of 1000 different measurements obtained in an acquisition time of six hours. The data points are taken in a random sequence for 31 magnetic field values equally distributed over the full range. Data points for 1 s and 3 s are recorded alternately. This way of data taking ensures that the signal is not influenced by residual long-term drifts of the experimental conditions.

The observed evaporation loss in Fig. 2 shows a pronounced dependence on the magnetic field which is to be compared with the expected cross section for elastic collisions as displayed in Fig. 1(b). After being very small at low magnetic fields, the loss increases for fields up to ~ 350 G where an expected local maximum of evaporative loss is observed. The loss then decreases and disappears at about 530 G as a consequence of the predicted zero crossing of the scattering length. Here the slight observed loss in the 1 s curve is explained by the finite ramp time of the magnetic field. In the 100 ms ramping time some evaporation does already take place. At 530 G the decrease of the trapped atom number between 1 s and 3 s is fully explained by rest gas losses without any further evaporation. For higher magnetic fields evaporative loss rapidly rises until it levels off at about 700 G. Up to the maximum attainable value of 1.5 kG high evaporation loss is observed. A slight decrease of the atom number for fields exceeding 1 kG occurs which we attribute to technical reasons; we observe an increasing noise for currents higher than ~ 130 A in the error signal of the resonator lock of the REDT. The relatively large and constant evaporative loss for fields exceeding 700 G is consistent with the predicted behavior of the cross section.

The evaporative cooling effect is confirmed by measuring the change of the truncation parameter η after 3 s of trapping at selected values of the magnetic field. For thermometry we turn off the magnetic field to avoid further elastic collisions and adiabatically lower the trap depth in a 1-s exponential ramp. The fraction of remaining atoms as a function of the relative depth then provides a good measure of η . At the zero-crossing at 530 G we observe only a slight increase of η to a value of 2.3(2) which is explained by the unavoidable evaporation during the magnetic field ramps. At 340 G close to the local maximum of $|a|$ we find an increase of η to 4.2(3) as a clear evidence of evaporative cooling. At 720 G, i.e. in the case of a large positive scattering length, a higher value of 5.5(2) is measured showing deeper evaporative cooling. Essentially the same η of 5.3(2) is obtained at $B = 1290$ G where scattering takes place in the triplet-dominated regime with a very large negative scattering




FIG. 3: Measurements on plain evaporation in magnetic fields close to the zero crossing of the scattering length under same conditions as the in Fig. 2 for a holding time of 3 s. Here the number of remaining atoms is normalized to the observed maximum value. The inset shows a series of measurements in a very narrow range around the maximum at 530(3) G together with a parabolic fit.

length.

The observed behavior is in agreement with a Monte-Carlo simulation taking into account the unitarity limitation of the cross section [19]. The results confirm the observed evaporation time scales and the attained values of η . We also find that collisions ejecting atoms out of the trap take place with a mean energy of $\sim 350 \mu\text{K}$ which corresponds to an effective wave number of $k \approx (300 a_0)^{-1}$.

We measure the minimum-loss feature in a closer range of magnetic fields to precisely determine the value of the magnetic field at which the zero crossing of scattering length occurs. The main data points in Fig. 3 are obtained with 500 individual measurements at a holding time of 3 s with the magnetic field randomly varied between 30 values in an interval between 370 G and 670 G; the data shown in the inset are obtained with 1000 measurements in the very narrow range between 520 G and 544 G. The results allow us to determine the B-field for minimum evaporative loss, and thus the zero-crossing of the scattering length to 530(3) G.

Forced evaporation measurements provide complementary data to plain evaporation and allow us to rule out a significant role of inelastic collisions. When the trap depth is ramped down, elastic collisions reduce trap loss in contrast to increased loss at constant trap depth. This can be understood by the spilling loss of energetic particles [18]: Without elastic collisions the most energetic particles are spilled out of the trap when its depth is reduced. With elastic collisions the evaporative cooling effect decreases the temperature and thus reduces the spilling loss.

In our forced evaporation measurements we reduce the trap depth in 10 s to 20% of its initial value in an exponential ramp and measure the number of remaining




FIG. 4: Fraction of atoms remaining after forced evaporation versus applied magnetic field. The trap depth is ramped down exponentially to 20% of the initial value in 10 s.

atoms; the results are displayed in Fig. 4. At reduced laser power of the REDT the magnetic field has to be restricted to 1 kG because of an increasing sensitivity of the resonator lock to current-dependent noise. A minimum number of atoms is now measured at 0 G and 530 G instead of the maximum observed with constant trap depth. The largest number of atoms is observed in the high-field region above 650 G as expected for the large scattering cross section.

In Ref. [20] inelastic collisions in ^6Li were studied at a temperature of $\sim 20\ \mu\text{K}$. A magnetic field dependent loss was observed around 680 Gauss, with evidence of an inelastic two-body process. To measure the loss rate under our conditions of higher temperature and much lower density we study the long-time evolution of samples which are pre-cooled by forced evaporation and then recompressed into a $750\ \mu\text{K}$ deep trap to a temperature of $\sim 150\ \mu\text{K}$ and a density of $5 \times 10^{10}\ \text{cm}^{-3}$. The magnetic field during the 60-s holding time is alternated for successive data points between 680 G and 300 G, the latter value corresponding to a very low loss rate in Ref. [20]. We find no significant difference in the number of remaining atoms, which leads to a clear upper bound to the two-body rate constant of $1 \times 10^{-12}\ \text{cm}^3/\text{s}$. This upper bound is lower than the value of $2 \times 10^{-12}\ \text{cm}^3/\text{s}$ of Ref. [20], whereas for a process involving higher partial waves one would expect the rate to increase with temperature. From our upper bound for inelastic loss and the measured values for η we infer that the ratio of good-to-bad collisions at very large positive and negative scattering length clearly exceeds 1000. Thus excellent conditions for evaporative cooling towards degeneracy can be expected near the Feshbach resonance.

In conclusion, our measurements confirm the predicted magnetic tunability of the s -wave scattering length in a spin-mixture of fermionic lithium atoms in the whole magnetic field range of experimental interest. The observed zero crossing of the scattering length at 530(3) G

together with the large cross section observed for higher fields provides clear evidence of the predicted Feshbach resonance. Moreover it enables more precise calculations of the ^6Li scattering properties. The resonance itself is masked by unitarity-limited scattering and requires much deeper evaporative cooling for a direct observation. The fact that we do not see any significant effect of inelastic loss highlights that the extremely large positive and negative scattering lengths attainable with fermionic lithium offer intriguing new possibilities for experiments on interacting Fermi gases.

Note added: Shortly before submission of the present manuscript we learned about measurements of the group of J.E. Thomas on the zero crossing of the scattering length which agree with our data.

We thank R.G. Hulet and H. Stoof for very useful discussions and V. Venturi for valuable input. Support by the Austrian Science Fund (FWF) within project P15115 and SFB 15 (project part 15) and by the Institut für Quanteninformation GesmbH is gratefully acknowledged.

-
- [1] S. Inouye *et al.*, Nature **392**, 151 (1998).
 - [2] P. Courteille *et al.*, Phys. Rev. Lett. **81**, 69 (1998).
 - [3] V. Vuletić, A. J. Kerman, C. Chin, and S. Chu, Phys. Rev. Lett. **82**, 1406 (1999).
 - [4] S. Cornish *et al.*, Phys. Rev. Lett. **85**, 1795 (2000).
 - [5] L. Khaykovich *et al.*, Science **296**, 1290 (2002).
 - [6] K. Strecker, G. Partridge, A. Truscott, and R. Hulet, Nature **417**, 150 (2002).
 - [7] T. Loftus *et al.*, Phys. Rev. Lett. **88**, 173201 (2002).
 - [8] S. Granade, M. Gehm, K. O'Hara, and J. Thomas, Phys. Rev. Lett. **88**, 120405 (2002).
 - [9] H. T. C. Stoof, M. Houbiers, C. Sackett, and R. Hulet, Phys. Rev. Lett. **76**, 10 (1996).
 - [10] M. Holland, S. Kokkelmans, M. Chiofalo, and R. Walser, Phys. Rev. Lett. **87**, 120406 (2001).
 - [11] E. A. Donley, N. R. Claussen, S. T. Thompson, and C. E. Wieman, Nature **417**, 529 (2002).
 - [12] D. Petrov, M. Holzmann, and G. Shlyapnikov, Phys. Rev. Lett. **84**, 2551 (2000).
 - [13] M. Houbiers, H. T. C. Stoof, W. McAlexander, and R. Hulet, Phys. Rev. A **57**, R1497 (1998).
 - [14] S. Kokkelmans *et al.*, Phys. Rev. A **65**, 053617 (2002).
 - [15] V. Venturi and C. Williams, private communication.
 - [16] A. Mosk *et al.*, Opt. Lett. **26**, 1837 (2001).
 - [17] C. Chin, V. Vuletić, A. J. Kerman, and S. Chu, Phys. Rev. Lett. **85**, 2717 (2000).
 - [18] O. Luiten, M. Reynolds, and J. T. M. Walraven, Phys. Rev. A **53**, 381 (1996).
 - [19] A. Mosk, unpublished.
 - [20] K. Dieckmann *et al.*, cond-mat/0207046.
 - [21] We use the analytical function $a(B) = (1 - \exp(-(B/p_1)^2)) \times (p_2(1+p_3/(B-p_4)) - p_5 \arctan(B-p_4))$ with $p_1 = 279$, $p_2 = -1528$, $p_3 = 202$, $p_4 = 816$, $p_5 = 312$ to approximate the calculated $a(B)$ of Refs. [13, 14, 15].